

HUMAN SACRAL HIATUS IN DRY BONES OF TELANGANA REGION & ITS CLINICAL SIGNIFICANCE DURING CAUDAL EPIDURAL ANESTHESIA – A MORPHOLOGICAL; MORPHOMETRICAL STUDY

Upendhar Reddy Pulluru¹, M. Venkateshwar Reddy *².

¹Tutor, Department of Anatomy, Sri Venkata Sai Medical College, Mahabubnagar, Telangana, India.

²Professor, Department of Anatomy, Sri Venkata Sai Medical College, Mahabubnagar, Telangana, India.

ABSTRACT

It is a very important to know the anatomical variations of sacral hiatus, for a successful epidural caudal anesthesia. Eleven direct morphometric measurements were done on sacral vertebrae and hiatus of 150 adult dry human sacral bones of unknown age and gender. Morphological features shape, level apex and base of SH were noted. Most common shape of SH was inverted inverted 'U' in 90 (60%) sacrum. In 62 (41.3%) sacra apex was present at the level of S4. The level of base of sacral hiatus – in 122(81.3%) at S5 level. Different mean values of 150 sacral hiatus are following. Height of sacral hiatus is 20.62mm; Width of hiatus is 12.11mm; distance from SH apex to S2 level is 29.38mm; distance from base of SH base to S2 level is 51.08mm; distance between upper border of S1 & SH apex is 60.08mm; depth of SH at apex level is 5.12mm; distance between two posterior superolateral crest is 60.20mm; distance between right posterior superolateral crest and apex of SH is 56.90mm; distance between leftpostero superolateral crest and apex of SH is 59.10mm; distance between right posterior superolateral crest and apex is 85.99mm; distance between left postero superolateral crest and apex of SH is 86.90mm.

KEY WORDS: sacral hiatus (SH), sacral canal, caudal epidural block (CEB).

Corresponding Author: Prof. Dr. M. Venkateshwarreddy, Professor of Anatomy, Sri Venkata Sai Medical College, Mahabubnagar, Telangana, India. **E-Mail:** dryemvee@gmail.com

Access this Article online	Journal Information
Quick Response code  DOI: 10.16965/ijar.2020.173	International Journal of Anatomy and Research RG Journal Impact: 0.21* ISSN (E) 2321-4287 ISSN (P) 2321-8967 https://www.ijmhr.org/ijar.htm DOI-Prefix: https://dx.doi.org/10.16965/ijar 
	Article Information
	Received: 04 Jun 2020 Peer Review: 06 Jun 2020 Revised: None
	Accepted: 24 Jun 2020 Published (O): 05 Jul 2020 Published (P): 05 Jul 2020

INTRODUCTION

Sicard and cathelin introduced caudal anesthesia in 1901, injected anesthetic drug through sacral hiatus into epidural space. In 1921 page and 1927 by dogliotti described lumbar epidural technique. Because of variations of sacral hiatus use of caudal epidural block declined. In caudal epidural block only cephalic distribution of the anesthetic agent occurs, so the loss of dermatomal segments are blocked[1]. Sacral canal is formed by sacral vertebral

foramina, triangular in section. Upper opening is seen on the basal surface. Its caudal opening is sacral hiatus. The dural and arachnoid sheaths of spinal cord terminates at the level of S2 vertebrae. The part of sacral canal below the S2 called caudal epidural space, which is filled with coccygeal nerves, venous plexus filum terminale and fats. This space is used for caudal epidural block for anesthesia and approach to it is through sacral hiatus[2]. Anatomical variations maybe shifting sacral

hiatus upwards or downwards, sometimes may absence of fusion of dorsal wall of sacrum (agenesis), and variations in shape, depth, narrowing, partial obliteration of sacral hiatus and canal, it may be traumatic or of pathological origin etc[3].

Sacral hiatus has been used for: 1) analgesia anesthesia in various clinical procedures, by injecting the drug into the epidural space through the hiatus in obstetrics & gynecology, hernia corrections, etc[4].

Sacral hiatal variations are always difficult for clinicians., 2) thecaloscopy[5], 3) trans-sacral endoscopy[6], 4) myelography[6], 5) caudal epidural injection of steroids / pain killers as a therapeutic agents[7] and 6) anterior epiduroscopic neural decompression: a minimal invasive spinal surgery[8].

So, the present study was conducted to find out the anatomical variations of sacral hiatus for caudal epidural block, with the help of morphometric measurements of the sacrum in relation to sacral hiatus in dry sacral bones.

MATERIALS AND METHODS

Fully ossified and un damaged 150 adult dry human sacral bones of unknown age and sex, obtained from the department of anatomy of Sri Venkata Sai Medical College And Hospital, Mahabubnagar, Telangana, India., were used. Measurements were taken by using a Vernier caliper of accuracy of 0.01mm, divider and steel measuring tape. The posterior closure and agenesis of dorsal wall bones were excluded. Morphological features, like shape, level of apex of sacral hiatus, level of base of sacral hiatus and following morphometric measurements were taken.

1. Height of sacral hiatus.
2. Width of sacral hiatus (distance between two sacral cornua).
3. Distance from the apex of sacral hiatus to the level of S2 foramen.
4. Distance from the base of sacral hiatus to the level of S2 foramen.
5. Distance between the upper border of S1 and sacral apex (length of sacral canal).
6. Depth of sacral hiatus at the level of its apex (anteroposterior diameter).

7. Distance between the two posterior superolateral crest.

8. Distance between right posterior superolateral crest and apex of sacral hiatus.

9. Distance between left posterior superolateral crest and apex of sacral hiatus.

10. Distance between right posterior superolateral crest to sacral apex.

11. Distance between left posterior superolateral crest to sacral apex.

Since the posterior superior iliac spine is readily palpable, impose on the superolateral sacral crest of sacrum. Line joins the both right and left superolateral crests of sacrum forms the base of triangle and the line joining the apex of sacral hiatus to both right and left superolateral crests, forms the other two arms of the triangle, which helps practically detect the triangle.

Statistical analysis: data were expressed as meancalculations and in percentages.

RESULTS

Agenesis of sacral hiatus in two sacrum, and complete agenesis of dorsal wall of sacrum found in one sacrum were excluded. Most common shape of hiatus was 'U' shaped found in 90(60%); in 42(28%) 'V' shaped; irregular shapes found in 8(5.3%); 'M' shaped found in 6(4%); dumbbell shaped in 4(2.6%) was out of total 150.

Apex was found in at the level of S4 in 62(41.3%) sacra. At the level of S3 are 32(21.3) && S2 are 6(4%).

Following different Mean values have found, height/length of sacral hiatus in 20.62mm; maximum width of hiatus found 12.11mm; distance from SH apex to S2 level 29.38mm; distance from SH base to S2 level 51.08mm; distance between upper border of S1 SH of apex 60.08mm; depth of SH at apex level 5.12mm; distance between two posterior superolateral crest 60.20mm; distance between right posterior superolateral crest and apex of SH 56.90mm; distance between left posterior superolateral crest and apex of SH 59.10mm; distance between right posterior superolateral crest and sacral apex 85.99mm; distance between left posterior superolateral crest and sacral apex 86.90mm.



Fig. 1: Completed agenesia of dorsal wall of Sacrum.



Fig. 5: Showing the "dumbbell" Shaped Sacral hiatus.



Fig. 2: Showing the "U" Shaped Sacral hiatus.

Table 1: Shapes of sacral hiatus.

S.No.	Shapes	Frequency (n)	%
1	Inverted 'U'	90	60
2	Inverted 'V'	42	28
3	Irregular	8	5.3
4	'M' shaped	6	4
5	Dumbbell	4	2.6
6	Total	150	100

Table 2: location of apex and base in relation to the sacral vertebrae.

S.No.	Level of vertebrae	Location of apex		Location of base	
		Frequency (n)	%	Frequency (n)	%
1	S5	-	-	122	81.3
2	S4	62	41.3	20	13.3
3	S3	32	21.3	-	-
4	S2	6	4	-	-
5	Coccyx	-	-	8	5.3



Fig. 3: Showing the "M" Shaped Sacral hiatus.

Table 3: Showing the results of different morphometric measurements.

S.No.	Parameters	Mean value
1	Height of Sacral Hiatus	20.62
2	Width of Sacral Hiatus	12.11
3	Distance from SH Apex to S2 level	29.38
4	Distance from SH Base to S2 level	51.08
5	Distance between Upper Border S1 & SH Apex	60.08
6	Depth of SH at apex level	5.12
7	Distance between two posterior superolateral crest	60.2
8	Distance between right posterior superolateral crest and apex of SH	56.9
9	Distance between left posterior superolateral crest and apex of SH	59.1
10	Distance between right posterior superolateral crest and sacral apex	85.99
11	Distance between left posterior superolateral crest and sacral apex	86.9



Fig. 4: Showing the "V" Shaped Sacral hiatus.



Fig. 6: Showing width measurement at the base of the Sacral hiatus.




Fig. 7: Showing width measurement at the base of the 2nd Sacral bone.



Fig. 8: Left posterior superolateral crest of apex.




Fig. 9: Showing measurement from left crest to apex of sacrum.



Fig. 10: Showing the measurement at upper border of S1 to apex of Sacral hiatus.




Fig. 11: Showing measurement of AP diameter of the Sacral hiatus.



Fig. 12: Showing measurement from Apex to S2 Level.



Fig. 13: Showing measurement of height of sacral hiatus.



Fig. 14: Showing measurement between two superolateral crest.



Fig. 15: Showing measurement of the Depth of Sacral hiatus at Apex.




Fig. 15: Showing measurement of the Depth of Sacral hiatus at Apex.

Table 4: Comparison of Shapes of Sacral Hiatus by different authors.

S.No.	Authors	Total number	Inverted'U' shape(%)	Inverted 'V' shape (%)	Irregular shape (%)
1	Osunwoke et al [14]	54	24.1	33.1	13
2	Vinod et al [15]		29.7	46.53	-
3	Nasr A Y et al [16]	150	31.33	38.66	15.33
4	Qudusia et al [13]	194	62.37	22.16	8.76
5	Deepa S et al [25]		57.5	25	17.5
6	Archana singh et al [33]	56	60.7	25	7.1
7	Present study	150	90	42	8

Table 5: Level of apex & base of sacral hiatus reported by different authors.

Authors	Total no. (N)	Level of Apex of Sacral Hiatus				Level of Base of Sacral Hiatus		
		S4	S3	S2	S5	S5	S4	Co
Nagar et al(2004) [17]	270	147(55.9%)	98(37.3%)	9(3.4%)	9(3.4%)	191(72.6%)	29(11.1%)	43(16.3%)
Dipali Rani et al(2012) [18]	160	80(50%)	72(45%)	—	8(5%)	132(82.5%)	14(8.7%)	14(8.7%)
Rammurti KS et al (2013) [4]	116	59(50.8%)	48(41.3%)	9(7.7%)	—	84(72.4%)	22(18.9%)	10(8.6%)
Nasr A Y et al (2014) [16]	150	81(54%)	22(14.66)	2(1.33%)	41(27.33%)	105(70%)	18(12%)	27(18%)
Archana singh et al (2015) [33]	56	28(50%)	24(42.9%)	4(7.1%)	—	46(82.1%)	8(14.3)	2(3.6)
Present study	150	70(46.6%)	66(44%)	8(5.3%)	6(4.3%)	128(85.3%)	18(14.3%)	4(2.6%)

Table 6: Height, Width and AP Diameters of sacral hiatus reported by different authors.

Authors	Total SH(mm)	Length of SH(mm)	Width of SH(mm)	AP diameter of SH(mm)
Senoglu et al (2005) [27]	96	32.1	17.47	4.46
Dipali Rani et al (2012) [18]	160	23.61 ± 8.28	12.75 ± 2.92	5.34 ± 1.39
Lakshmi et al (2013) [31]	51	34.96 ± 12.9	14.8 ± 2.32	4.61 ± 1.5
Sanatnu B et al (2013) [32]	100	35.92 ± 3.75	9.79 ± 1.31	7.23 ± .71
Clarista MQ et al (2013) [19]	104	24.73 ± 9.38	16.87 ± 3.66	5.58 ± 1.39
Nasr A Y et al (2014) [16]	150	27.16 ± 1.29	11.50 ± 3.13	4.78 ± 1.68
Mishra M et al (2014) [21]	93	19.73	12.11	4
Archana singh et al (2015) [33]	56	21.73 ± 8.92	11.59 ± 3.25	5.02 ± 2.09
Present study	150	24.52 ± 8.32	13.65 ± 1.42	5.28 ± 1.44

Table 7: Comparison of Depth (Diameter) of Sacral hiatus at its Apex.

Authors	Total no. of sacrum	Mean diameter (mm)
Blanchis et al [29] (by USG)	-	5.3 mm
Chen et al [11] (by USG)	-	6 mm
Crigton et al [30] (by MRI)	15	4.6 mm
Archana singh et al [33]	56	5.02 ± 2.09 mm
Present study	150	5.06 mm

Table 8:Distance from apex of SH to S2 foramen (C).

Authors	Total Number	Mean distance (C) (mm) ± SD
Rammurti KS et al [4]	116	30.2 ± 10.5
Dipali Rani et al [18]	160	31.33 ± 10.59
Lakshmi et al [31]	—	31.53 ± 9.05
Mustafa MS et al [23]	46	41 ± 11.4
Santanu et al [32]	100	43.41 ± 8.5
Archana singh et al [33]	56	30.30 ± 11.01
Present study	150	38.40 ± 10.6

Table 9: Triangle between two Post Superolateral Crestand apex of sacral hiatus (EFG) reported by different authors

Authors (N)	Base (E) (mm) +SD	Right margin (F) (mm) + SD	Left margin (G) (mm) + SD	Type of triangle
Senoglu et al [27](2005)n=96	66.51 ± 53.52	67.10 ± 9.95	67.53 ± 9.48	equilateral triangle
Lakshmi et al [31] (2013)n=51	61.51 ± 19.49	58.28 ± 9.36	58.54 ± 9.47	equilateral triangle
Sanatnu B et al [32](2013) n=100	64.77 ± 5.2 mm	58.41 ± 2.70	58.37 ± 2.76	isosceles triangle
Aggarwal et al [28] (2009)n= 114	50.96 ± 6.69	59.92 ± 8.84	59.99 ± 8.31	Equilateral triangle -45% 55% cases both sides were much shorter than the base of the triangle.
Rammurti K S et al [4] (2013)n=116	69.5 ± 5.8	61.4 ± 11.2	57.4 ± 9.7	Isosceles triangle
S Deepa et al [25](2014)n=40	62.3 ± 4.4	69 ± 14.5	71 ± 17.7	Isosceles triangle
Archana singh et al [33] (2015)n=56	61.16 ± 5.42	57.54 ± 10.2	58.32 ± 10.59	equilateral triangle
Present study n=150	64.6 ± 8.48	57.24 ± 6.8	62.22 ± 8.6	equilateral triangle

DISCUSSION

For successful sacral caudal epidural block, the knowledge of anatomical variations and structural modifications of sacral hiatus is necessary. Caudal epidural block is a technique of injecting medication in to epidural space via sacral hiatus. It is practiced since 1952 to anaesthetize lumbar and sacral dermatomes and

also for the symptomatic relief of low backache disorders in patients by injecting corticosteroids. Injecting technique used is prone position of patient, table is flexed or with pillow beneath the pelvis, followed by palpation of sacral cornu and hiatus with the help of left hand fingers. Needle is then inserted 2-3 mm at an angle of 45 degrees, 'pops' through the sacrococcygeal ligament[3]. A failure rate has been reported by

some authors; in children 14.82% failure rate noted by Paolo Busoni et al[9] and in adults Shu-Yam Wong[10] reported a highest success rate 95.9%. Ultrasound guided needle placement success was 100% by Carl P. C. Chen[11]. The apex of the sacral hiatus is an important bony point for caudal epidural block but there may be difficulty in its palpation in some patients especially in obese. Hence other bony landmarks can also be used in relation with sacral hiatus. We can use a triangle which is drawn between the posterior superior iliac spines (superolateral sacral crest in dry sacral bones) and the apex of sacral hiatus. We measured the distance between these points which can guide the clinician for detection of sacral hiatus easily and increase the success rate of caudal epidural block. William R. Meeker[12] and colleagues noted that the diameter of sacral canal is 4.5mm and it is difficult to pass the needle into the sacral hiatus if it cannot be located accurately.

In present study the shape of sacral hiatus were variable and the most common being the inverted 'U' shaped in 90(60%) sacrum, followed by inverted 'V' shaped in 14(25%) sacra and this was similar to Qudusia et al[13] (2014) in which they found the most common shape of sacral hiatus was the inverted 'U' shape in 121(62.37%) sacra and inverted 'V' shaped in 43(22.16%) sacra but was not similar with the Osunokietal[14] and Vinod et al[15]. Vinod et al (1992) noted most common shape of sacral hiatus was inverted 'V' shape in 94(46.55%) sacra and inverted 'U' shape 58(38.66%) sacra and inverted 'U' shape in 47(31.33%) Egyptian sacrum.

In present study 4(2.6%) sacra was found to have its outline like dumbbell which was very low when compared with other studies as by Nagaret al[17](2004) in 36(13.3%) sacra and Dipali Rani et al[18](2012) in 20(12.5%) sacra. M shaped sacral hiatus was observed in 6(4%) sacra in this study which was near to the study done by Clarista MQ et al[19] (2013) noted in 3(2.9%) sacra.

In present study apex of the SH was most commonly found at the level of S₄ sacral vertebrae, in 62(41.3%), which was almost similar with the observations by Nagar et al [17] (2004) in 147 (55.9%), by Dipali Rani et al[18](2012) in 80(50%)

by Rammurti KS et al[4] (2013) in 59(50.8%) sacra and by Nasr A Y et al[16] (2014) in 81(54%) sacra. The base of sacral hiatus was seen most commonly at the level of S5 vertebrae, in 122(81.3%) sacra in present study similar with the Dipali Rani et al[18](2012) in 132(82.5%) ,Manisha B Sinha et al[20](2014) in 55(88.7%) sacra and by Mishra M et al[21] (2014) in 79(89.77%)sacra.

Cornua are palpable as a landmark on either side just below the apex of hiatus for localization of the sacral hiatus clinically. In the present study the distance between the two cornu was found to be 12.11mm. Sekiguchi et al[26] (2004) noted 10.2mm, Senoglu et al[27](2005) noted 17.47mm, Aggarwal et al[28] (2009) noted 11.95mm, Blanchais et al[29](2010) noted 14.2mm, Nasr A Y et al[16] (2014) noted 11.5mm, Mishra M et al[21] (2014) noted 12.11mm and S Deepa et al[25] (2014) noted 12.2mm.

Diameter(anterio-posterior length of SH at the apex) of sacral hiatus is important because if it is <2mm, then there may be difficulty in use of 22 G needle for Caudal epidural space. In present study 12.5% sacra had AP diameter <2mm and lowest measured value was 1.64mm and the mean value was 5.12mm (given in table4). Similar observation were noted by Kumar et al[15] (1992) 4.8mm, by Nagar et al[17](2004) 4.8mm, by Sekiguchi et al[26](2004) 6mm, by Senoglu et al[27](2005)4.46mm, by Mishra M et al[21](2014) 4mm and by Nasr A Y et al[16](2014) 4.78±1.87mm in Egyptian sacra. AP diameter by ultrasonographic study was measured by Blanchisetal[29] and Chen et al[11] and found to be mean AP diameter 5.3mm and 6mm respectively. By MRI study done by Crigtonetal [30] AP diameter found to be 4.6 mm, which was similar to the present study.

Length of sacral canal is measured as vertical distance between the apex of SH and the upper border of the sacral canal; in present study it is found to be 60.08mm which is similar to that measured by Mourgella et al[5] using MRI 59.03mm. Patil et al[3](2012) noted 64.77±17.07 mm, Lakshmi et al[31] (2013) noted 60.41±13mm and Santanu et al[32](2013) noted 66.19±4.8mm. As the dural sac ends at the level of S₂ vertebrae, the distance from the SH apex to S2 spine

and distance from the base of SH to apex of S2 spine noted, because anaesthetist or endoscopist must know that how far to push the needle or instrument into the sacral canal from the SH. In present study the distance between apex of SH to spine of S2 was found to be mean of 29.38mm. By other authors it was reported as Rammurti KS et al [4] (2013) 30.2±10.5mm, by Dipali Rani et al [18] (2012) 31.33±10.59 mm and by Lakshmi et al [31] (2013) 31.53±9.05mm which was similar with the present study. The distance between the base of SH and S2 spine was reported in present study was 51.08 mm and by other authors it was found to be as by Rammurti K S et. al [4] (2013) 59.5±8.8mm, by Dipali Rani et al [18] (2012) 54.88±7.92mm which was similar with the present study but by Santanu et al [32] (2013) and Phalgunan et al [24] (2013) it was noted as 79.64±8.9mm and 60.23±17.97mm respectively which was little higher than present study.

A triangle formed between the two posterior superior iliac spines and the apex of SH, used as an important landmark to locate the SH clinically. In this study this triangle had almost equilateral features. The mean value of three sides of triangle i.e. base, right margin and left margin were 61.16±5.42mm, 57.54±10.2mm and 58.32±10.59mm respectively, these values are nearly equal and formed an equilateral triangle which is similar to other studies done by Senoglu et al [27] (2005) mean values of three sides were 66.51±53.52mm, 67.10±9.95mm and 67.53±9.48 mm

Respectively and forming equilateral triangle. By Lakshmi et al [31] (2013) the values were 62.6±10.48mm, 56.24±6.8mm and 64.22±8.6mm respectively, forming almost equilateral triangle. By Mustafa MS et al [23] (2012) these values were 75.5±10.3mm, 75±10.2mm and 75±10.2mm, forming equilateral triangle, but by some authors it was not found to be equilateral triangle as by Sanatnu B et al [32] (2013) it was an isosceles triangle and mean values were 64.77±5.2mm, 58.41±2.70mm and 58.37±2.76 mm and by Aggarwal et al [28] (2009) reported mean values of three sides were 50.96±6.69mm, 59.92±8.84mm and 59.99±8.31mm respectively and found equilateral triangle only in 45% cases and in rest 55% cases both sides were much

shorter than the base of the triangle.

CONCLUSION

There are variabilities in the anatomical structure of SH. This may be due to genetic and racial factors. The given landmarks in this study, resulting the formation of equilateral triangle, which can provide the practical benefit to the clinician for localization of SH in CEB. Further clinical trials are required to compare the existing techniques and our anatomical description to support the result of this study.

Conflicts of Interests: None

REFERENCES

- [1]. Cap Steven J Zito. Adult Caudal Anaesthesia: A Reexamination of the Technique. *Journal of the American Association of Nurse Anaesthesia*. 1993;61(2):153-157.
- [2]. Susan Standring. *The Anatomical Basis of Clinical Practice*. Gray's Anatomy, Elsevier Churchill Livingstone, London. 40th edition: 724-28, 760-61.
- [3]. Patil Dhananjay S, Jadav H R, Binod K, Mehta C D, Patel Vipul D. Anatomical Study of Sacral Hiatus for Caudal Epidural Block. *National Jor of Medical Research*. 2012;2(3):272-75.
- [4]. Rammurti K S, Anil K Reddy Y. Anatomical Study of Sacral Hiatus for Successful Caudal Epidural Block. *Int J Med Res Health Sci*. 2013;2(3):496-500. <https://doi.org/10.5958/j.2319-5886.2.3.086>
- [5]. Mourgela S, Sakellaropoulos A, Anagnostopoulou S, Warnke J P. The Dimension of sacral Spinal Canal in Thecaloscopy: A Morphometric MRI Study. *Neuroanatomy*. 2009;8:1-3.
- [6]. Sheila B Jones, Shaw D W W, Lawrence E Jacpbson. A Transsacral Approach Through the Sacral Hiatus for Myelography. *AJR*. 1997;169:1179-1181. <https://doi.org/10.2214/ajr.169.4.9308486> PMid:9308486
- [7]. Bentley A. Caudal Epidural Steroid Injections. *Pain Physician*. 2000;3(3):305-312.
- [8]. Eirch O Richter, Abramova M V, Canta F, Andres J D, Peter L, Manchiaro P et al. Anterior Epiduroscopic Neural Decompression: Eight Center Experiences in 154 Patients. *European J of Pain Supplements*. 2011;5:401-407. <https://doi.org/10.1016/j.eujps.2011.08.006>
- [9]. Paolo Busoni, Armando Sarti. Sacral Intervertebral Epidural Block. *Anesthesiology*. 1987;67:993-995. <https://doi.org/10.1097/00000542-198712000-00023> PMid:3688545
- [10]. Shu-Yam Wong et al., Caudal Epidural Block for Minor Gynecologic Procedures in Outpatient Surgery. *Chang Gung Med J*. 2004;27(2):116-121.
- [11]. Carl P. C. Chen et al. Ultrasound Guidance in Caudal Epidural Needle Placement. *Anesthesiology*. 2004;101:181-4. <https://doi.org/10.1097/00000542-200407000-00028> PMid:15220789

- [12]. William R. Meeker. Sacral Nerve Block Anesthesia: The anatomy involved, Technique and Clinical Application. *Annals of Surgery*. 1924;80(5):739-772. <https://doi.org/10.1097/00000658-192411010-00012> PMID:17865134 PMCID:PMC1399815
- [13]. Qudusia S, Shariff MH, Jacob M, Rao CP, Avadhani R. A Morphological study of Sacral Hiatus with its Clinical Implications. *Indian Journal of Applied Research*. 2014;4(2):34-7. <https://doi.org/10.15373/2249555X/FEB2014/135>
- [14]. Osunwoke E, Oladipo G, Allison TA, Orlu E. A Study of Sacral Hiatus in Dry Human Sacra in Southern Nigeria. *Journal of Biology, Agriculture and Healthcare*. 2014; 4(5):43-8.
- [15]. Kumar V, Pandey SN, Bajpai RN, Jain PN, Longia GS. Morphometric Study of Sacral Hiatus. *JASI* 1992;41(1):7-13.
- [16]. Naser A Y, Ali Y H, Naser A E. The Sacral Hiatus: An Anatomical Study of Both Cadaveric and Dry bones. *Translational Clinical Biology*. 2014; 2(1):4-12. <https://doi.org/10.14259/tcb.v2i1.124>
- [17]. Nagar SK. A Study of Sacral Hiatus in Dry Human Sacra. *J. Anat. Soc. India*. 2004;53(2):18-21.
- [18]. Dipali Rani Pal, Rahman M A, Fatema K. Morphometric Study of Sacral Hiatus: A Basis for Successful Caudal Epidural Block. *Bangladesh J of Anatomy*. 2012;10(1):5-10. <https://doi.org/10.3329/bja.v10i1.15750>
- [19]. Clarista MQ, Gautham K. Morphometrical Study of Sacral Hiatus In Dry Human Sacra in West Indian Population. *Journal of Surgery* 2013;2(2):56-63.
- [20]. Manisha B Sinha, Rathor M, Sinha H P. A Study of Variation of Sacral Hiatus in Dry Bone in Central Indian Region. *International Jour of Healthcare and Biomedical Research*. 2014;2(4):46-52.
- [21]. Mishra M, Singh A K, Jha S, Satyavathi P, Sah N. Sacral Hiatus in Dry Human Sacra. *Janki Medical College Journal of Medical sciences*. 2014;2(1):17-22. <https://doi.org/10.3126/jmcjms.v2i1.11391>
- [22]. Patel ZK, Thummar B, Rathod SP, Singel TC, Patel S, Zalawadia A. Multi-centric Morphometric Study of Dry Human Sacrum of Indian Population in Gujarat Region. *NJIRM*. 2011;2(2):31-35.
- [23]. Mohamed SM, Omayma MM, El Raouf HHA, Hosam MA. Morphometric Study of Sacral Hiatus in Adult Human Egyptian Sacra: Their Significance in Caudal Epidural Anesthesia. *Saudi Journal of Anaesthesia*. 2012;6(4):350-7. <https://doi.org/10.4103/1658-354X.105862> PMID: 23493625 PMCID:PMC3591553
- [24]. Vijisha Phalgunan, Baskaran S. Morphometrical Analysis of Sacral Hiatus and its Clinical Significance. *The Health Agenda*. 2013;1(1):10-15.
- [25]. Deepa S, Rajesekar S S. Anatomical Study of Sacral Hiatus. *Int J of healthcare and Bioresearch*. 2014;3(1):31-35.
- [26]. Sekiguchi M, Yabuki S, Saton K, Kikuchi S. An anatomical Study of the Sacral Hiatus: A Basis for Successful Caudal Epidural Block. *Clin J Pain*. 2004;20:51-4. <https://doi.org/10.1097/00002508-200401000-00010> PMID:14668657
- [27]. Senoglu N, Senoglu M, Oksuz H, Gumusalan Y, Yuksel KZ, Zencirci B et al. Landmarks of the Sacral Hiatus for Caudal Epidural Block: An Anatomical Study. *British Journal of Anaesthesia*. 2005;95(5):692-5. <https://doi.org/10.1093/bja/aei236> PMID:16155035
- [28]. Aggarwal A, Aggarwal A, Harjeet, Sahni D. Morphometry of Sacral Hiatus and its Clinical Relevance in Caudal Epidural Block. *Surgical and Radiological Anatomy*. 2009;31(10):793-800. <https://doi.org/10.1007/s00276-009-0529-4> PMID:19578805
- [29]. Blanchais A, Le Goff B, Guillot P, Berthelot JM, Glemarec J, Maugars Y. Feasibility and Safety of Ultrasound - guided Caudal Epidural Glucocorticoid Injections. *Joint Bone Spine*. 2010;77(5):440-444. <https://doi.org/10.1016/j.jbspin.2010.04.016> PMID:20869897
- [30]. Crighton IM, Barry BP, Hobbs GJ. A Study of the Anatomy of the Caudal Space Using Magnetic Resonance Imaging. *Br J Anaesth*. 1997;78(4):391-395. <https://doi.org/10.1093/bja/78.4.391> PMID:9135359
- [31]. Lakshmi Trikkur Anantharaman, Azra Jabeen. Surgically Relevant Morphometry of Sacral Hiatus. *Anatomica Karnataka*. 2013;7(1):52-56.
- [32]. Santanu B, Majumdar S, Chakraborty P, Majumdar Sibanu, Mazumdar A. A Morphometric Study of Sacral Hiatus for Caudal Epidural Block among the Population of West Bengal. *Indian Journal of Basic and Applied Medical Research*. 2013;7(2):660-67.
- [33]. A Singh, Natl J Integr, Morphological and Morphometrical Study of Sacral Hiatus of Human Sacrum *NJIRM* 2018; 9(4):65-73.

How to cite this article:

Upendhar Reddy Pulluru, M. Venkateshwar Reddy. HUMAN SACRAL HIATUS IN DRY BONES OF TELANGANA REGION & ITS CLINICAL SIGNIFICANCE DURING CAUDAL EPIDURAL ANESTHESIA – A MORPHOLOGICAL; MORPHOMETRICAL STUDY. *Int J Anat Res* 2020;8(3.1):7609-7616. DOI: 10.16965/ijar.2020.173