

Case Report

ACCESSORY RENAL ARTERY AND ITS CLINICAL SIGNIFICANCE

Sunday Abba *, Amanuel Tesfay, Peter Ekanem.

Department of Human Anatomy College of Health Sciences Mekelle University, Ethiopia.

ABSTRACT

Knowledge of the existence of the aberrant renal arteries is important because they may be inadvertently damaged during renal surgery and their presence must be considered in evaluating a donor kidney for possible renal transplantation. Using conventional dissecting techniques, the posterior abdominal wall was dissected in a 45years old embalmed male cadaver in the gross anatomy dissecting laboratory of college of health Sciences Ayder Campus Mekelle University Ethiopia.

Following the fine dissection, we noticed an aberrant right renal artery (accessory renal artery) originated from the abdominal aorta at the level of L3 vertebra, besides the normal renal artery that arises high up at the level of L1, L2. This accessory renal artery does not have a parallel course with the normal renal artery, it course upward and laterally to the inferior pole of the right kidney which it pierces, However there was no such observation on the left.

KEYWORDS: Accessory Renal Artery; Abdominal Aorta; Arterial abnormalities; Renal Transplant.

Address for Correspondence: Sunday Abba. Department of Human Anatomy College of Health Sciences Mekelle University, P.O.BOX 1974. Mekelle, Ethiopia. **E-Mail:** suneabba007@yahoo.co.uk

Access this Article online

Quick Response code



Web site: International Journal of Anatomy and Research
ISSN 2321-4287
www.ijmhr.org/ijar.htm

Received: 13 Nov 2013

Peer Review: 13 Nov 2013 Published (O):30 Dec 2013

Accepted: 28 Dec 2013 Published (P):30 Dec 2013

INTRODUCTION

The increasing use of invasive diagnostic and interventional procedures in cardiovascular diseases makes it imperative that the type and frequency of vascular variations be well documented and understood. With the advent of laparoscopic renal surgeries, and donor nephrectomies, it becomes necessary for surgeons to understand the variations that exist in the renal vasculature otherwise renal laparoscopic and nephrectomies could be jeopardized [1].

The renal arteries are paired branches of the abdominal aorta giving out at the level of L1, L2. The paired renal arteries take 20% of cardiac output to supply organs that represent less than one-hundredth of total body weight. They supply the kidneys through a number of subdivisions described sequentially as segmental, lobar, interlobar, and arcuate arteries.

These are end arteries with no anastomoses. The arcuate arteries further divide into interlobular arteries which give rise to the afferent arteries to the glomeruli.

The renal arteries branch laterally from the aorta just below the origin of the superior mesenteric artery. Both cross the corresponding crus of the diaphragm at right angles to the aorta.

The right renal artery is longer and often higher, passing posterior to the inferior vena cava, right renal vein, head of the pancreas and descending part of the duodenum. The left renal artery is a little lower and passes behind the left renal vein, the body of the pancreas and splenic vein. It may be crossed anteriorly by the inferior mesenteric vein.

A single renal artery to each kidney is present in 70% of individuals. The arteries vary in their level of origin and in their calibre, obliquity and precise relations.

In its extrarenal course each renal artery gives off one or more inferior suprarenal arteries, a branch to the ureter and branches which supply perinephric tissue, the renal capsule and the pelvis. Near the renal hilum, each artery divides into an anterior and a posterior division, and these divide into segmental arteries supplying the renal vascular segments. Accessory renal arteries are common (30% of individuals), and usually arise from the aorta above or below the main renal artery and follow it to the renal hilum. They are regarded as persistent embryonic lateral splanchnic arteries. Accessory vessels to the inferior pole cross anterior to the ureter and may, by obstructing the ureter, cause hydronephrosis. Rarely, accessory renal arteries arise from the coeliac or superior mesenteric arteries near the aortic bifurcation or from the common iliac arteries.

Existence of aberrant arteries is accountable in cases of renal pathologies, radiological interventions, renal transplants and other surgical approach on them. Altered state of hemodynamics was thought of in cases of multiple arteries supplying the renal [2].

In this present study we reported the unusual branching of the accessory right renal artery giving out by the aorta at the level of L3 vertebra and its associated clinical significance.

CASE REPORT

Using conventional dissecting techniques, the posterior abdominal wall was dissected in a 45years old embalmed male cadaver in the gross anatomy dissecting laboratory of college of health Sciences Ayder Campus Mekelle University Ethiopia. The medical history of this cadaver was not available. Following the fine dissection, we noticed an aberrant right renal artery (accessory renal artery) originated from the abdominal aorta at the level of L3 vertebra, besides the normal renal artery that arises high up at the level of L1, L2. This accessory renal artery does not have a parallel course with the normal renal artery, it course upward and laterally to the inferior pole of the right kidney which it pierces as shown in figure1. This accessory renal artery arises from a different position as reported by Ramesh Rao in 2010. He reported the presence of aberrant renal arteries that aroused high up and very close to the renal artery. There was no such observation on the left.



a- Abdominal Artery, b- Renal Artery, c- Accessory Renal Artery, d- Right Ureter, e- Right Testicular Vessels

Fig. 1: Showing Accessory Renal Artery.

DISCUSSION

Knowledge of the existence of the aberrant renal arteries is important because they may be inadvertently damaged during renal surgery and their presence must be considered in evaluating a donor kidney for possible renal transplantation. Persistence of certain of the cephalic mesonephros vessels, however may result in the arterial abnormalities [3].

These mesonephric arteries extend from C6 – L3 during development. Most cranial vessels disappear while the caudal arteries persist to form a network, the rete arteriosum urogenitale that supply the future metanephros. The metanephros in future develop into the adult kidney deriving its blood supply from the from the lowest suprarenal artery which gives out a permanent renal artery. Persistent roots of the network form these segmental arteries of the adult kidneys having variations at their point of origin. The kidney grafts with multiple arteries resulted in post transplant morbidity and graft loss following the ligation of the polar arteries. The transplantation of the kidneys with the single renal artery is technically easier compared to the kidneys with multiple arteries [4].

Aberrant or accessory renal arteries have been of interest to the clinicians for some years mainly because of the possible part the vessel play in the causation of hydronephrosis. However judging by the many description of these vessels in the literature, it is evident that there is no established criterion for aberrance: the term have been applied equally to an additional artery

in the renal pedicle, or to a vessel entering the kidney at either pole whether derived from the main artery or from the aorta or a branch of the aorta [5].

Literatures have shown that aberrant renal arteries are common in fused kidneys but this was not the case here. Aberrant arteries perforate the substance of the kidney rather than entering its Hilum as observed here, the accessory renal artery pierced the inferior pole of the kidney. These arteries could arise as high as the inferior phrenic arteries or as low as the internal iliac arteries. The unusual vessel may originate from the aorta, as well as gonadal, common iliac, median sacral artery, external and internal iliac arteries, superior or inferior mesenteric arteries [6].

The presence of additional renal artery is very probable when the main renal artery has a diameter of less than 4.15mm. Here we measure the diameter of the main right renal artery and it was found to be 3.87mm. Kidney presenting a main renal artery diameter to be 5.5mm very probably do not present with additional renal arteries. So the renal artery diameter is a factor which should be considered as predicting the presence of additional renal arteries [7].

CONCLUSION

Considering the increase in incidences of the accessory renal arteries and their clinical implications in renal laparoscopic surgery, hypertension, renal transplantations and hydronephrosis, it becomes necessary to report such observation any where it is found so that surgeon will become more aware of this occurrence. Otherwise, renal transplant and renal laparoscopy may be jeopardized. This is also important for academic and radiological procedures.

Conflicts of Interests: None

REFERENCES

- [1]. T. Ramesh Rao. Aberrant renal arteries and its clinical implication: a case report. International journal of anatomical variations. 2011 40:37-39
- [2]. Standring Sed. Grays anatomy. The anatomical basis of clinical practice. 40th Ed. Edinburg Churchill and Livingstone. 2008: 1231,1233
- [3]. Cerny JC, Karsch D. Aberrant renal arteries. Urology. 1973; 2: 623–626.
- [4]. Ozkan U, Oguzkurt L, Tercan F, Kizilkilic O, Koc Z, Koca N. Renal artery origins and variations: Angiographic evaluation of 855 consecutive patients. Diagn Interv Radiol. 2006; 12: 183–186.

- [5]. Graves FT. The aberrant renal artery. J Anat. 1956; 90: 553–558.
- [6]. Bergman RA, Thomson SA, Afifi AK, Saadeh FA. Compendium of Human Anatomic Variation. Baltimore, Urban & Schwarzenberg. 1988; 81–83.
- [7]. Saldarriaga B, Pinto SA, Ballesteros LE. Morphological expression of the renal artery. A direct anatomical study in Columbian half-caste population. Int J Morphol. 2008; 26; 31–38.

How to cite this article:

Sunday Abba, Amanuel Tesfay, Peter Ekanem. ACCESSORY RENAL ARTERY AND ITS CLINICAL SIGNIFICANCE. Int J Anat Res 2013;03:155-57.